



UltraSal-ER™

(28.5% Salicylic Acid Extended Release)
Antiviral Film-Forming Solution

Patient Instructions

Your healthcare professional has prescribed UltraSal-ER, a topical prescription preparation for the treatment of common and plantar warts. In order for UltraSal-ER to work correctly and to ensure maximum benefit, the following instructions should be followed carefully. Of course, as with any medication, always consult your healthcare professional if you experience any discomfort or unexpected reactions.

STEP 1: WASH



- Soak wart area in warm water for about 5 minutes.
- Remove any loose tissue using a wash cloth, emery board or pumice stone.
- Dry thoroughly.

STEP 2: BRUSH



- Using the specially designed brush for precise application, apply twice to entire wart surface, allowing the first application to dry before applying the second.
- Be careful not to apply to surrounding skin.
- Once applied, a protective, antiviral water-resistant seal forms.

Apply as directed by your healthcare professional.

Clinically visible improvement will normally occur during the first or second week of therapy.

UltraSal-ER is indicated for the topical treatment and removal of common warts and plantar warts.

Important Safety Information

For topical use only. Not for ophthalmic, oral, or intravaginal use. A localized irritant reaction may occur if UltraSal-ER is applied to the normal skin surrounding the wart. Patients with diabetes or impaired blood circulation should not use UltraSal-ER. UltraSal-ER should not be used on moles, birthmarks, and unusual warts with hair growing from them, or warts on the face.

See full Prescribing Information for UltraSal-ER including Important Safety Information at www.UltraSalER.com